arXiv:1409.6890v1 [math.CA] 24 Sep 2014

Supersolutions to degenerated logistic equation type

Marcos Marvá.
Departamento de Física y Matemáticas, Universidad de Alcalá,
28871 Alcalá de Henares, Spain
marcos.marva@uah.es

Mathematics Subject Classification (2010): 35J99 (primary).
Keywords: Supersolution, logistic equation.

Abstract

In this work we provide a method for building up a strictly positive supersolution for the steady state of a degenerated logistic equation type, i.e., when the weight function vanishes on the boundary of the domain. This degenerated system is related in obtaining the so-called large solutions. Previously, this problem was handled as the limit case of non degenerated approaching problems. Our method can be adapted straightforwardly to degenerated boundary value problems.

1 Introduction

In this work we show how to build up a positive strict supersolution to the boundary value problem

$$\begin{cases} -\Delta u = \lambda m(x)u - a(x)f(x,u) & in \quad \Omega, \\ u = g(x) & on \quad \partial\Omega, \end{cases} \tag{1}$$

where $a \in C^{\mu}(\bar{\Omega};[0,+\infty))$, $m \in C^{\mu}(\bar{\Omega};\mathbb{R})$, $\mu \in (0,1)$, $a \equiv 0$ on $\partial\Omega$, $g \in C^{1+\mu}(\partial\Omega;[0,+\infty))$ is such that $g \geq 0$, $g \not\equiv 0$, $f \in C^{\mu,1+\mu}(\bar{\Omega}\times[0,+\infty);\mathbb{R})$, $f(x,0) = 0$, $f(x,u) > 0$, $\partial_u f(x,u) > 0$ and $\lim_{k\nearrow+\infty} \frac{f(x,ku)}{k} = +\infty$ for each $u > 0$ uniformly in $x \in \Omega$. The set $\Omega \subset \mathbb{R}^N$ is a bounded domain with $\partial\Omega$ of class C^2.

Problem (1) (with $m \equiv 1$) stands for the steady states of the generalized logistic growth law [13]. Typically, u stands for the distribution in Ω of individuals of certain species, λ stands for the net growth rate of u modulated by $m(x)$, $a(x)$ simulates demography pressure and, along with $m(x)$, environmental heterogeneity. When $a \equiv 0$ somewhere, model (1) (with $m \equiv 1$) becomes a Malthusian model in the region $\Omega_0 := \{x \in \Omega;\ a(x) = 0\}$, known as *refuge*. We deal with the case $a \equiv 0$ on $\partial\Omega$, but it could be adapted to apply in the general case when $\Omega_0 \neq \emptyset$ and $\Omega_0 \subset \Omega$.

In addition, solving problem (1) is an intermediate step to prove the existence of large solutions related to problem (1), i.e., a function $u \in C^{2,\nu}(\overline{\Omega})$ such that

$$\begin{cases} -\Delta u = \lambda m(x)u - a(x)f(x,u)u & in \ \ \Omega, \\ \lim\limits_{x\to\partial\Omega} u(x) = \infty, \end{cases} \tag{2}$$

see [11]. Large solutions come across from studies concerning combustion due to Keller [9] and Osserman [14]. In fact, there are available uniqueness results for problem (2) ([3], [5], [12] and references therein). As long as we know, problem (1) has been handled as the limit case of non degenerated approaching problems.

If we restrict ourselves in problem (1) to $a(x) \geq \gamma > 0$ on $\overline{\Omega}$, it is easy to show that there exists $\epsilon > 0$ small enough such that $\epsilon\varphi$ is a positive subsolution to problem (1), being φ the principal eigenfunction of $-\Delta$ in Ω under homogeneous Dirichlet boundary conditions. In this case a positive constant K such that

$$K > \max\left\{ \max_{x\in\partial\Omega}\{g\}\,,\ \frac{\max\limits_{x\in\overline{\Omega}}\{\lambda m(x)\}}{\min\limits_{x\in\overline{\Omega}}\{a(x)\}} \right\}, \tag{3}$$

provides us with a supersolution to problem (1). Now, enlarging K if necessary, we get an ordered pair $(\epsilon\varphi, K)$ of sub-supersolution. Thanks to a Theorem by Amann [1], there exists a solution to problem (1) between $\epsilon\varphi$ and K. Unfortunately, as soon as we let $a = 0$ somewhere in $\bar{\Omega}$, we lose condition (3), so it is needed something different from a constant to get a supersolution to problem (1). Different strategies have been used to avoid this problem: some authors approximate Ω by $\Omega_n := \{x \in \Omega;\ \mathrm{dist}(x, \partial\Omega) > 1/n\}$ (for each n $a(x)$ is uniformly bounded away from 0 in Ω_n, that is $a \geq \gamma > 0$, see [4] and [11]), and others approach $a(x)$ by means of $\frac{1}{n} + a(x)$, essentially in order to avoid condition $a \equiv 0$ on $\partial\Omega$ and to generate a sequence of solutions of approximate problems converging to a solution to problem (1), see [2] and [8].

2 Results

The result we present here allows us to obtain via sub and supersolution the existence of a solution to (1). Actually, the construction of the supersolution in Ω follows from an extension of the Faber [6] and Krahn [7] inequality due to López-Gómez [10] which provides us with a lower estimate for the main eigenvalue to $-\Delta$ on the domain D:

$$\sigma_1[-\Delta, D] \geq \frac{\sigma_1[-\Delta, B_1(0)] \cdot |B_1(0)|^{2/N}}{|D|^{2/N}}, \tag{4}$$

where $B_1(0) := \{x \in \mathbb{R}^N; |x| \leq 1\}$, $\sigma_1[-\Delta, B_1(0)]$ is the principal eigenvalue of $-\Delta$ on $B_1(0)$ and $|D|$ stands for the Lebesgue's measure of D.

Theorem 2.1 *Consider the boundary value problem* (1)*. Then, for each* $\lambda \in \mathbb{R}$*, there exists an ordered pair* $(\underline{u}, \overline{u})$ *consisting of a subsolution and a positive strict supersolution, where* $\underline{u}(x) < \overline{u}(x)\ \forall x \in \Omega$.

Proof.- Let us define

$$O_\epsilon := \{x \in \mathbb{R}^N;\ \mathrm{dist}(x, \partial\Omega) < \epsilon\}, \qquad \sigma_\epsilon := \sigma[-\Delta, O_\epsilon],$$

where, keeping in mind (4), for a fixed $\lambda \in \mathbb{R}$ we can choose $\epsilon > 0$ such that

$$\max_{x \in \overline{\Omega}} \{\lambda m(x)\} < \sigma_\epsilon.$$

Let φ_o^ϵ be the principal eigenfunction of $-\Delta$ under homogeneous boundary conditions on O_ϵ. We define

$$\Phi := \begin{cases} \varphi_o^\epsilon & \text{in } \overline{\Omega} \cap O_{\epsilon/2}, \\ \Psi & \text{in } \Omega_{\epsilon/2} := \Omega \backslash (\overline{\Omega} \cap O_{\epsilon/2}), \end{cases}$$

where $\Psi \geq \tau > 0$ is any regular enough function such that $\Phi \in C^2(\overline{\Omega})$. We are looking for the existence of a constant $K > 0$, large enough, such that function

$$\overline{u} := K\Phi$$

is a strict positive supersolution to problem (1).

We estimate now K on $\overline{\Omega} \cap O_{\epsilon/2}$. Keeping in mind that $a(x) \geq 0$ and $K\Phi = K\varphi_o^\epsilon$, it must be

$$\begin{cases} K(-\Delta\varphi_o^\epsilon) \geq \lambda m K\varphi_o^\epsilon - af(x, K\varphi_o^\epsilon) & \text{in } \overline{\Omega} \cap O_{\epsilon/2}, \\ K\varphi_o^\epsilon \geq g > 0 & \text{on } \partial\overline{\Omega} \cap O_{\epsilon/2} = \partial\Omega. \end{cases}$$

In the interior of the domain it should happen

$$K(-\Delta\varphi_o^\epsilon) \geq \lambda m K\varphi_o^\epsilon - af(x, K\varphi_o^\epsilon),$$

which is equivalent to

$$K\sigma_\epsilon\varphi_o^\epsilon \geq \lambda m K\varphi_o^\epsilon - af(x, K\varphi_o^\epsilon)$$

by the definition of φ_o^ϵ. Rearranging terms we get

$$af(x, K\varphi_o^\epsilon) \geq K\varphi_o^\epsilon \left(\lambda m - \sigma_\epsilon\right), \tag{5}$$

where the left hand side is non negative and the right one is strictly negative because of the selection of ϵ. Therefore, (5) is a proper inequality $\forall K > 0$. By construction, on the boundary we have $\varphi_o^\epsilon|_{\partial\Omega}, g > 0$ and therefore, whenever

$$K > \max_{x \in \partial\Omega} \left\{ \frac{g(x)}{\varphi_o^\epsilon(x)} \right\}, \tag{6}$$

the boundary condition is satisfied (with strict inequality).

In the domain $\Omega_{\epsilon/2}$ it should be verified

$$K(-\Delta\Psi) \geq \lambda m K\Psi - af(x, K\Psi). \tag{7}$$

rearranging terms, expression (7) is equivalent to

$$\frac{f(x,K\Psi)}{K} \geq \max_{x\in\overline{\Omega}_{\epsilon/2}} \left\{ \frac{\Delta\Psi + \lambda m\Psi}{a} \right\}. \tag{8}$$

As $a(x) > 0 \quad \forall x \in \Omega_{\epsilon/2}$, function $\frac{\Delta\Psi+\lambda m\Psi}{a}$ reaches its maximum (provided a regular Ψ). By hypothesis on f, there exists K_0 such that $\forall K > K_0$ condition (8) holds. Thus, there exists $K > 0$ such that function $K\Phi$ is a strict positive supersolution to (1). Function $\underline{u} := 0$ is a subsolution to problem (1), and $\underline{u}(x) < \overline{u}(x)\ \forall x \in \bar{\Omega}$. ■

Corollary 2.2 *For each $\lambda \in \mathbb{R}$, problem* (1) *has an unique positive solution.*

Proof.- Since we have an ordered pair formed by a subsolution and a supersolution, a Theorem due to Amann [1] guaranties the existence of a solution u to problem (1) such that $0 \leq u \leq K\Psi$. Uniqueness follows, for instance, from [11]. ■

Acknowledgement.- The author thanks E. Gómez-Hidalgo for her fruitful comments and support. The author is partially supported by Ministerio de Ciencia e Innovación (Spain), projects MTM2011-24321 and MTM2011-25238.

References

[1] H. Amann, *Fixed point equations and nonlinear eigenvalue problems in ordered Banach spaces*, SIAM Rev., **18**:219 (1976), 620–709.

[2] F.C. Cirstea, V. Radulescu, *Asymptotics for the blow-up boundary solution of the logistic equation with absorption*, C. R. Math. Acad. Sci. Paris, **226** (3)336 (2003), 231–236.

[3] F.C.Cirstea, V. Radulescu, *Nonlinear problems with boundary blow-up: a Karamata regular variation theory approach*, Asymptot. Anal., **46**:3–4 (2006), 275–298.

[4] Y. Du, Q. Huang, *Blow-up solutions for a class of semilinear elliptic and parabolic problems*, SIAM J. Math. Anal., **31**:1 (1999), 1–18, .

[5] Y. Du, *Asymptotic behavior and uniqueness results for boundary blow-up solutions*, Differ. Integral Equ. Appl., **17** (2004), 819–834.

[6] G. Faber *Beweis das unter allen homogenen Membranen von Gleicher Fläche und gleicher Spannung die kreisförmige den tiefsten Grundton gibt*, Munch. Ber. (1923), 169–172.

[7] E. Krahn, *Über eine von Rayleigh formulierte Minimaleigenschaft des Kreises*, Math. Ann., **94** (1925), 97–100.

[8] J. García-Melián, R. Letelier-Albornoz, J.C. Sabina de Lis, *Uniqueness and asymptotic behaviour for solutions of semilinear problems with boundary blow-up*, Proc. Amer. Math. Soc., **129**:12 (2001), 3593–3602.

[9] J.B. Keller, *On solutions of* $\Delta u = f(u)$, Comm. Pure Appl. Math., **10** (1957), 503–510.

[10] J. López-Gómez, *The maximum principle and the existence of principal eigenvalues for some linear weighted boundary value problems*, J. Diff. Eqns. Conf., **1**, No.1 electronic only (2000), (1996) 263–294.

[11] J. López-Gómez, *Large solutions, metasolutions, and asymptotic behaviour of the regular positive solutions of sublinear parabolic problems*, J. Diff. Eqns. Conf., **5** (2000), 135–171.

[12] J. López-Gómez, *Optimal uniqueness theorems and exact blow-up rates of large solutions*, J. Diff. Eqns., **224**:2 (2006), 385–439.

[13] J.D. Murray, *Mathematical biology.* Berlin: Biomathematics 19 Springer-Verlag, 1993.

[14] R. Osserman, *On the inequality* $\Delta u \geq f(u)$, Pacific. J. Math., **7** (1957), 1641–1647.